

National Suicide Prevention Lifeline: After an Attempt



**A Guide for Medical
Providers in the
Emergency Department
Taking Care of Suicide
Attempt Survivors**



U.S. DEPARTMENT OF HEALTH AND HUMAN SERVICES
Substance Abuse and Mental Health Services Administration
www.samhsa.gov

NATIONAL
SUICIDE
PREVENTION
LIFELINE™
1-800-273-TALK

RED
para la
PREVENCIÓN
de
SUICIDIO
NACIONAL
1-888-628-9454

If you are in crisis, please call:

National Suicide Prevention Lifeline
1-800-273-TALK (8255)

En español: 1-888-628-9454

TTY: 1-800-799-4889

www.suicidepreventionlifeline.org



National Suicide Prevention Lifeline: *After an Attempt*

**A Guide for Medical Providers in the
Emergency Department Taking Care of
Suicide Attempt Survivors**

Acknowledgements

This brochure was originally developed by the National Alliance on Mental Illness (www.nami.org) in partnership with the Suicide Prevention Resource Center (www.sprc.org) under Grant Number 1.U79 SM55029-01 from the Substance Abuse and Mental Health Services Administration (SAMHSA), U. S. Department of Health and Human Services (HHS). The views, policies, and opinions expressed are those of the authors and do not necessarily reflect those of SAMHSA or HHS.

Public Domain Notice

All material appearing in this publication is in the public domain and may be reproduced or copied without permission from SAMHSA or the Center for Mental Health Services. Citation of the source is appreciated. This publication may not be reproduced or distributed for a fee without the specific, written authorization of the Office of Communications, SAMHSA, HHS.

Electronic Access and Copies of Publication

This publication can be accessed electronically through www.mentalhealth.samhsa.gov. For additional free copies of this document, please contact SAMHSA's National Mental Health Information Center at www.mentalhealth.samhsa.gov, 1-800-789-2647, or 1-866-889-2647 (TDD). Ask for Publication No. CMHS-SVP-0161.

Recommended Citation

U.S. Department of Health and Human Services.
*National Suicide Prevention Lifeline:
After an Attempt. A Guide for Medical Providers in the
Emergency Department Taking Care of Suicide Attempt
Survivors.* CMHS-SVP-0161, Rockville, MD: Center
for Mental Health Services, Substance Abuse and
Mental Health Services Administration, 2006.

Originating Office

Substance Abuse and Mental Health
Services Administration
1 Choke Cherry Road
Rockville, MD 20857

CMHS-SVP-0161

Printed 2006

SUICIDE: Helping Patients and Their Families

After an Attempt

The emergency department (ED) is the frontline of medicine and often serves as the doorway into the medical system for people in distress, including people who attempt suicide. Individuals who have attempted suicide are at increased risk for later dying by suicide, and up to 20 percent of those who attempted suicide in the past will try again in the future.¹ The ED is often the initial point of contact with the health system for many of these individuals, and it offers a unique opportunity to help people who have attempted suicide to begin to recover from the depression and hopelessness that led to their suicide attempt.

The purpose of this brochure is to provide you with some quick tips to enhance care in the ED for people who have attempted suicide, while also providing information on the Health Insurance Portability and Accountability Act (HIPAA), patient discharge, and resources about suicide for medical professionals, patients, and their families.

¹ Appleby L., et al. Suicide Within 12 Months of Contact With Mental Health Services: National Clinical Survey, *British Medical Journal* 318(7193):1235-1239, May 1999.

Patient Care in the Emergency Department: Helpful Tips

Medical and psychological assessment and the development of a suicide prevention or safety plan are key steps in treating an individual who has attempted suicide. Your medical training no doubt provided you with the tools to perform a thorough assessment of an individual who has attempted suicide, including exploring physical conditions that may contribute to a person's behavior. However, for more information on this topic, the American Psychiatric Association's *Practice Guideline for the Assessment and Treatment of Patients With Suicidal Behaviors* is an excellent resource (www.psych.org/psych_pract/treatg/pg/pg_suicidalbehaviors.pdf or call 703-907-7300).

In addition to your assessment, communicating with others who may have more information about a patient's history, such as a family member or another physician, often can provide valuable insight as you assess the patient's situation. Any such communications about a patient's protected health information must be in accordance with the Federal patient privacy standards known as the HIPAA. Here are some brief highlights of HIPAA:

Communicating With a Patient's Family or Other Caregiver

- HIPAA allows the disclosure of protected health information to a family member or caregiver in the following situations:
 - If the patient consents to his or her information being shared.
 - If hospital personnel provide the patient with the opportunity to object to the disclosure, and the patient does not express an objection.
 - If it is reasonable to infer from the circumstances that the patient does not object to the disclosure.
- If an emergency physician, based on his or her best medical judgment, feels the patient lacks the capacity to provide informed consent, and that disclosure is in the patient's best interest, then the emergency physician can communicate directly with a patient's family or other caregiver, even if a patient objects.
- However, information shared with family members or other caregivers should be limited to that which is directly relevant to the caregiver's involvement with the patient's care or payment for that care.

Remember: Although confidentiality laws may restrict you from communicating medical information to nonmedical attendants, HIPAA allows you to receive unlimited information from families or other caregivers without patient consent.

Communicating With Other Medical Professionals About a Patient

If possible, consult with an individual's inpatient and/or outpatient provider to help maintain continuity of care and allow for informed decisionmaking by the individual and his or her treatment team.

As long as a good faith effort has been made to inform your patient about your institution's privacy practices, HIPAA allows you to:

- Share information about the person with other medical providers who are involved in the person's care, both within and outside of your own institution. This applies to all forms of communication (e.g. verbal, electronic, written).
- Provide your name and contact information to another medical provider for communication about the person.

To read more about HIPAA online, visit www.cms.hhs.gov/hipaa.

Patient Discharge From the Emergency Department: What the ED Can Do To Ease the Transition

In addition to your assessment and interventions for a person in the ED, a well-conceived discharge plan will go a long way in helping people safely and successfully begin to recover. Here are a few things to consider when developing a person's discharge plan.

Before leaving the ED, it is recommended that the patient and his or her family, if appropriate, should have:

- An understanding of discharge arrangements.
- A written statement with information about prescribed medication(s) and treatment plans.
- Key contacts to call—including outpatient providers, crisis lines, and peer-support centers.
- Specific instructions about the signs, symptoms, or conditions that require a return.

Other key points to discuss with patients *and* families before discharge include:

- What to look for that may indicate a return of suicidal feelings.
- Followup care—Who? Where? When?
- How to get resources and supports in the community.
- How to reduce the immediate hazards of another suicide attempt (such as information on removing or restricting items that are frequently used for self-harm).
- Who to call with questions or concerns.

If a patient is being discharged and an appointment for followup care cannot be arranged before discharge, strongly encourage the patient to seek followup care within the first few days of returning home. In areas where this service is available, mobile crisis teams have been effective in helping patients in crisis connect with outpatient treatment following an emergency visit. Consider linking your patient with a team in your area to increase the chances that he or she will access followup care.

Also remind your patient that the emergency department is open 24 hours a day, 365 days a year to help, if he or she continues to have thoughts of suicide or if the medical team is unavailable to provide the needed care.

One of the most important things you can do for a patient or family member after having been in the ED is to offer hope. Patients and families will look to you to determine the prognosis and for some assurance that this will not happen again. While you cannot guarantee there will not be a recurrence, you *can* assure them that recovery is likely if the individual and, if appropriate, a family member work closely with a therapist to ensure that the safety and treatment plans are meaningful and effective.

The following pages of this brochure offer a list of crisis lines and referrals for more information on suicide and mental illness

for patients and their families. Please also consider providing your patient with a copy of the brochure *Suicide: Taking Care of Yourself After an Attempt*, and its companion brochure for families, *Suicide: Taking Care of Family Members After an Attempt*. Each brochure includes information on safety and recovery, as well as additional resources for help.

Following these tips should increase the likelihood that your patients who have attempted suicide find the longer-term care that will prevent another attempt—one that could be fatal. You and the professional team around you can make the difference.

Resources for Professionals in the Emergency Department

The following resources offer excellent information on suicide and patient assessment.

American Association of Suicidology

A resource and education organization dedicated to the understanding and prevention of suicide. For more information, visit www.suicidology.org or call 202-237-2280.

American College of Emergency Physicians (ACEP)

A national medical society committed to advancing emergency care through continuing education, research, and public education. For more information, visit www.acep.org or call 1-800-798-1822.

American Foundation for Suicide Prevention (AFSP)

The AFSP has recently produced an excellent poster for emergency physicians on the evaluation of patients who attempt suicide; to obtain a free copy, visit www.afsp.org or call 1-888-333-AFSP.

American Psychiatric Association *Practice Guideline for the Assessment and Treatment of Patients with Suicidal Behaviors*

Visit www.psych.org/psych_pract/treatg/pg/pg_suicidalbehaviors.pdf or call 703-907-7300. Gliatto, M. and Rai, A., Evaluation and Treatment of Patients with Suicidal Ideation. *American Family Physician*, March 15, 1999. Full text available online at www.aafp.org/afp/990315ap/1500.html.

National Strategy for Suicide Prevention

A comprehensive national plan that lays a framework for action to prevent suicide. For more information, visit www.mentalhealth.samhsa.gov/suicideprevention.

Suicide Prevention Resource Center

Supports suicide prevention with the best of science, skills, and practice. For more information, visit www.sprc.org or call 1-877-GET-SPRC (438-7772).

Resources for Patients and Families in the Emergency Department

In a crisis, contact:

1-800-273-TALK (8255)

TTY: 1-800-799-4TTY (4889)

National Suicide Prevention Lifeline

A 24-hour, toll-free crisis hotline that links callers to a nearby crisis center. The Lifeline accepts calls from non-English speakers.

www.suicidepreventionlifeline.org

*For more information about suicide and
mental illness:*

American Association of Suicidology

A resource and education organization dedicated to the understanding and prevention of suicide.

www.suicidology.org or call 202-237-2280

American Foundation for Suicide Prevention

Dedicated to advancing the public's knowledge of suicide and its prevention.

www.afsp.org or call 1-888-333-AFSP

Befrienders International/Samaritans

An online resource that gives support through e-mail and offers a directory of local crisis helplines.

www.befrienders.org

**Covenant House Neline Hotline
1-800-999-9999**

A 24-hour, toll-free crisis hotline offering confidential and immediate crisis intervention and referrals to community resources.

www.covenanthouse.org/programs_nl.html

**Hispanic Community Resource
Helpline
1-800-473-3003**

(La Linea Nacional de Ayuda)

Offers support for Latinos who need information about educational, health, and human service providers.

**Link's National Resource Center for
Suicide Prevention and Aftercare
(LINK-NRC)**

Provides suicide-related community education in prevention, intervention, aftercare, and support.

www.thelink.org or call 404-256-9797

**National Alliance on Mental Illness
(NAMI)**

Offers information, support, and advocacy for persons affected by mental illnesses.

www.nami.org or call 1-800-950-NAMI
(6264)

National Disability Rights Network

Serves individuals with a wide range of disabilities by guarding against abuse, advocating for basic rights, and ensuring system accountability. The Web site provides a directory of member agencies by State.

www.ndrn.org or call 202-408-9514 or 202-408-9521 (TTY)

National Institute of Mental Health (NIMH)

The leading Federal agency for research on mental and behavioral disorders.

www.nimh.nih.gov or call 1-866-615-6464

National Mental Health Association

Addresses all aspects of mental health and mental illness.

www.nmha.org or call 1-800-969-NMHA (6642)

National Mental Health Information Center (NMHIC)

A SAMHSA-operated Center that provides information about mental health and is available on weekdays from 8:30 a.m. to 12 a.m. Eastern Standard Time to answer mental health questions.

www.mentalhealth.samhsa.gov or call 1-800-789-2647 or 1-866-889-2647 (TDD)

National Organization for People of Color Against Suicide

Addresses and raises awareness about suicide in minority communities.

www.nopcas.com or call 1-866-899-5317

Suicide Awareness Voices of Education (SAVE)

Dedicated to preventing suicide through education, public awareness, and stigma reduction.

www.save.org or call 952-946-7998

Suicide Prevention Action Network (SPAN) USA

A national organization dedicated to action and advocacy for suicide prevention.

www.spanusa.org or call 202-449-3600

Suicide Prevention Resource Center

Supports suicide prevention with the best of science, skills, and practice.

www.sprc.org or call 1-877-GET-SPRC (438-7772)

The Trevor Helpline

1-866-4U-TREVOR

A national 24-hour, toll-free suicide prevention hotline aimed at gay and questioning youth.

www.thetrevorproject.org

These resources may contain materials that express views, policies, and opinions that do not necessarily reflect those of the Substance Abuse and Mental Health Services Administration and the U.S. Department of Health and Human Services.

Your Notes Here:

Your Notes Here:

the fact that the *Journal of Applied Behavior Analysis* is the most widely read journal in the field of behavior analysis.

It is my hope that this book will be useful to you in your current or future work.

Thank you for your interest in this book. I am sure you will find it a valuable addition to your library.

Very truly yours,
John M. Hayes

John M. Hayes, Ph.D., is a professor of psychology at the University of California, Santa Barbara. He is also a past president of the American Psychological Association.

He has published numerous articles and books on the psychology of language and the psychology of education. He is also a frequent speaker at national and international conferences.

Dr. Hayes is currently working on a book on the psychology of writing. He is also working on a book on the psychology of reading. He is also working on a book on the psychology of learning.

He is also working on a book on the psychology of memory. He is also working on a book on the psychology of intelligence. He is also working on a book on the psychology of personality.

He is also working on a book on the psychology of social behavior. He is also working on a book on the psychology of development. He is also working on a book on the psychology of aging.

He is also working on a book on the psychology of health. He is also working on a book on the psychology of law. He is also working on a book on the psychology of religion.

He is also working on a book on the psychology of art. He is also working on a book on the psychology of music. He is also working on a book on the psychology of sports.

He is also working on a book on the psychology of politics. He is also working on a book on the psychology of economics. He is also working on a book on the psychology of history.

He is also working on a book on the psychology of philosophy. He is also working on a book on the psychology of science. He is also working on a book on the psychology of technology.

He is also working on a book on the psychology of culture. He is also working on a book on the psychology of society. He is also working on a book on the psychology of the individual.

He is also working on a book on the psychology of the mind. He is also working on a book on the psychology of the brain. He is also working on a book on the psychology of the body.

He is also working on a book on the psychology of the soul. He is also working on a book on the psychology of the spirit. He is also working on a book on the psychology of the heart.

He is also working on a book on the psychology of the hand. He is also working on a book on the psychology of the foot. He is also working on a book on the psychology of the eye.

He is also working on a book on the psychology of the ear. He is also working on a book on the psychology of the nose. He is also working on a book on the psychology of the mouth.

He is also working on a book on the psychology of the tongue. He is also working on a book on the psychology of the throat. He is also working on a book on the psychology of the chest.

He is also working on a book on the psychology of the stomach. He is also working on a book on the psychology of the intestines. He is also working on a book on the psychology of the bladder.

He is also working on a book on the psychology of the rectum. He is also working on a book on the psychology of the vagina. He is also working on a book on the psychology of the penis.

He is also working on a book on the psychology of the testis. He is also working on a book on the psychology of the ovary. He is also working on a book on the psychology of the uterus.

He is also working on a book on the psychology of the fallopian tube. He is also working on a book on the psychology of the placenta. He is also working on a book on the psychology of the fetus.

He is also working on a book on the psychology of the newborn. He is also working on a book on the psychology of the infant. He is also working on a book on the psychology of the child.

He is also working on a book on the psychology of the adolescent. He is also working on a book on the psychology of the young adult. He is also working on a book on the psychology of the adult.

For More Information:



American College of
Emergency Physicians®

ADVANCING EMERGENCY CARE 



The Nation's Voice on Mental Illness

www.nami.org

Printed 2006 • CMHS-SVP-0161